



A one-parameter deformation of the Farahat-Higman algebra

Jean-Paul Bultel

► To cite this version:

Jean-Paul Bultel. A one-parameter deformation of the Farahat-Higman algebra. European Journal of Combinatorics, 2011, pp.308-323. 10.1016/j.ejc.2010.10.008 . hal-00825448

HAL Id: hal-00825448

<https://hal.science/hal-00825448>

Submitted on 23 May 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A ONE-PARAMETER DEFORMATION OF THE FARAHAT-HIGMAN ALGEBRA

JEAN-PAUL BULTEL

ABSTRACT. We show, by introducing an appropriate basis, that a one-parameter family of Hopf algebras introduced by Foissy [Adv. Math. 218 (2008) 136-162] interpolates between the Faà di Bruno algebra and the Farahat-Higman algebra. Its structure constants in this basis are deformation of the top connection coefficients, for which we obtain analogues of Macdonald's formulas.

1. INTRODUCTION

The center Z_n of the group algebra $\mathbb{Z}[\mathfrak{S}_n]$ of the symmetric group is spanned by conjugacy classes C_μ , which are parametrized by partitions μ of n . The *connection coefficients* $a_{\mu\nu}^\lambda$ are the structure constants

$$(1) \quad C_\mu C_\nu = \sum_{\lambda \vdash n} a_{\mu\nu}^\lambda C_\lambda$$

of Z_n . These coefficients, whose calculation is in general very hard, have important applications to various enumerative problems or to the calculation of certain matrix integrals [7].

For a partition $\mu = (\mu_1 \geq \mu_2 \geq \dots \geq \mu_r > 0)$, define

$$(2) \quad \bar{\mu} = (\mu_1 - 1, \mu_2 - 1, \dots, \mu_r - 1),$$

the *reduced cycle type* of any permutation of cycle type μ . Denoting by $c_\rho(n)$ the conjugacy class C_μ of \mathfrak{S}_n such that $\bar{\mu} = \rho$, we can rewrite (1) in the form

$$(3) \quad c_\mu(n) c_\nu(n) = \sum_{\lambda} a_{\mu\nu}^\lambda(n) c_\lambda(n).$$

It has been proved by Farahat and Higman [4] that the connection coefficients $a_{\mu\nu}^\lambda(n)$ are polynomial functions of n , and are independent of n if $|\lambda| = |\mu| + |\nu|$. These are called the *top connection coefficients*.

One may use the top connection coefficients to define an algebra R , spanned by formal symbols c_μ indexed by all partitions, with multiplication rule

$$(4) \quad c_\mu c_\nu = \sum_{|\lambda|=|\mu|+|\nu|} a_{\mu\nu}^\lambda c_\lambda.$$

This is the Farahat-Higman algebra ([4], see also [10, ex. 24 p 131]). It is also proved in [4] that R is isomorphic to the algebra of symmetric functions Λ . The construction of an explicit isomorphism $\varphi : \Lambda \rightarrow R$ is more recent, and due to Macdonald [10, ex.

25 p 132] (another proof has been given by Goulden and Jackson [6]). More precisely, Macdonald constructed a basis (g_μ) of Λ such that

$$(5) \quad g_\mu g_\nu = \sum_{|\lambda|=|\mu|+|\nu|} a_{\mu\nu}^\lambda g_\lambda,$$

and obtained a recursive formula for the calculation of $a_{\mu\nu}^\lambda$.

Murray [12] explicitated the images in Λ of the projections of symmetric functions of Jucys-Murphy elements in R under this isomorphism. In Section 3, we give a new derivation of his result, and new proofs of various results of Biane [1] and Matsumoto-Novak [11].

It is well-known that the algebra of symmetric functions is a Hopf algebra. Its standard coproduct, denoted here by Δ_0 , comes from its interpretation as the algebra of polynomial functions on the multiplicative group G_0 of formal power series with constant term 1. It has, however, another Hopf algebra structure, known as the *Faà di Bruno algebra*, coming from its interpretation as the algebra of polynomial functions on the group $G_1 = \{ta(t) | a \in G_0\}$ of formal diffeomorphisms of the line under composition (see, *e.g.*, [3]). In fact, Macdonald's basis g_μ is the dual of the image $h_\mu = S_1(h_\mu)$ of the basis of complete homogeneous functions h_μ by the antipode of the Faà di Bruno algebra \mathcal{F} .

This suggests to interpret g_μ as living in the dual \mathcal{F}^* of \mathcal{F} . However, contrary to R , this dual is not commutative, being the universal enveloping algebra of the Lie algebra \mathfrak{g}_1 of G_1 . Thus, such an interpretation does not make sense *a priori*.

To clarify this situation, we make use of a one-parameter deformation \mathcal{F}_γ of \mathcal{F} recently discovered by Foissy [5] in his investigation of combinatorial Dyson-Schwinger equations in the Connes-Kreimer algebra. We then obtain for the structure constants of \mathcal{F}_γ in the dual basis of $S_1(h_\mu)$ a one-parameter deformation of Macdonald's formulas which are recovered for $\gamma = 0$.

We follow the conventions of [10]. For the convenience of the reader, the most essential ones are recalled in Section 2.

2. NOTATIONS AND BACKGROUND

2.1. Partitions. Let n be a positive integer. A finite sequence of strictly positive integers $(\lambda_1, \lambda_2, \dots)$ is called a *partition* of n if $\lambda_1 \geq \lambda_2 \geq \dots$ and $\lambda_1 + \lambda_2 + \dots = n$. We then write $\lambda \vdash n$. The λ_i are the *parts* of λ , $|\lambda| = n$ is the *weight* of λ and the number $l(\lambda)$ of parts in λ is the *length* of λ . The *multiplicity* $m_\lambda(k)$ of k in λ is the number of parts in λ equal to k . We set

$$(6) \quad z_\lambda = \prod_{k \geq 1} k^{m_k(\lambda)} m_k(\lambda)!$$

2.2. The algebra of symmetric functions. We denote by Λ the algebra of symmetric functions. The bases $(m_\lambda), (e_\lambda), (h_\lambda), (p_\lambda)$ and (s_λ) are respectively the *monomial*, *elementary*, *complete*, *power sum* and *Schur* symmetric functions. These bases of Λ are parametrized by partitions of all integers. For any basis (b_λ) , we denote by b_n

the symmetric function $b_{(n)}$. Denote by $\langle \cdot, \cdot \rangle$ the usual *scalar product* on Λ , for which (s_λ) is an orthonormal basis. For this scalar product, (p_λ) is an orthogonal basis, and one has

$$(7) \quad \langle p_\lambda, p_\mu \rangle = \delta_{\lambda\mu} z_\mu$$

The bases (m_λ) and (h_λ) are dual to each other.

$$(8) \quad \langle h_\lambda, m_\mu \rangle = \delta_{\lambda\mu}$$

(δ is the Kronecker symbol). We denote by p_n^\perp the *adjoint* of the multiplication by p_n , in the sense

$$(9) \quad \langle p_n f, g \rangle = \langle f, p_n^\perp(g) \rangle$$

where f and g are any two symmetric functions. This operator is a *derivation*. More precisely,

$$(10) \quad p_n^\perp = n \frac{\partial}{\partial p_n}$$

This operator acts on the h_n as follows

$$(11) \quad p_n^\perp h_N = h_{N-n}$$

for any $N > n$, so that

$$(12) \quad p_n^\perp = \sum_{r \geq 0} h_r \frac{\partial}{\partial h_{n+r}}$$

Now, let X and Y be two alphabets. For any symmetric function f and any scalar k , we identify f with $f(X)$, and we define the algebra morphisms $f \rightarrow f(X + Y)$, $f \rightarrow f(kX)$ and $f \rightarrow f(k)$ by

$$(13) \quad p_n(X + Y) = p_n(X) + p_n(Y)$$

$$(14) \quad p_n(kX) = k p_n(X)$$

$$(15) \quad p_n(k) = k$$

The standard Hopf algebra structure of Λ is defined by the coproduct Δ_0

$$(16) \quad \Delta_0(f) = f(X + Y)$$

where we identify $f(X + Y)$ with an element of $\Lambda \otimes \Lambda$ by identifying $f \otimes g$ with $f(X)g(Y)$. The counit ϵ and the antipode S_0 are given by

$$(17) \quad \epsilon(f) = f(0)$$

and

$$(18) \quad S_0(f) = f(-X)$$

Denote by \mathcal{H}_0 this Hopf algebra.

$$(19) \quad \mathcal{H}_0 = (\Lambda, \cdot, 1, \Delta_0, \epsilon, S_0)$$

One can give another interpretation of \mathcal{H}_0 . Let

$$(20) \quad G_0 = \{a \mid a(t) = 1 + a_1 t + a_2 t^2 + \dots\} = 1 + t\mathbb{C}[[t]]$$

be the multiplicative group of formal power series with constant term 1, and let \mathcal{H} be the algebra of polynomial functions on G_0 . Let $k_n \in \mathcal{H}$ be the map defined by

$$(21) \quad k_n(a) = a_n$$

The standard coproduct for functions on a group is

$$(22) \quad \Delta f(a, b) = f(ab)$$

where Δf is identified with an element of $\mathcal{H} \otimes \mathcal{H}$. Then, $h_n \mapsto k_n$ is an isomorphism of Hopf algebras $\mathcal{H}_0 \rightarrow \mathcal{H}$.

2.3. Other bases of the algebra of symmetric functions. The formal series

$$(23) \quad u = tH(t) = t + h_1 t^2 + h_2 t^3 + \dots$$

has an inverse for the composition, so that we can rewrite t in terms of u , as follows.

$$(24) \quad t = u + h_1^* u^2 + h_2^* u^3 + \dots$$

The h_k^* are homogeneous symmetric functions of degree k , and the algebra morphism ψ from Λ to Λ defined by $\psi(h_k) = h_k^*$ is an *involution* :

$$(25) \quad \psi^2 = Id_\Lambda$$

The Lagrange inversion formula shows that $(n+1)h_n^*$ is the coefficient of t^n in $H(t)^{-(n+1)}$, so that

$$(26) \quad h_n^* = \frac{h_n(-(n+1)X)}{n+1}$$

We can define a multiplicative \mathbb{Z} -basis (h_λ^*) of Λ , by

$$(27) \quad h_\lambda^* = h_{\lambda_1}^* h_{\lambda_2}^* \dots = \psi(h_\lambda)$$

In [10, ex. 24 p 35], Macdonald shows that

$$(28) \quad (n+1)h_n^* = \sum_{\lambda \vdash n} (-1)^{l(\lambda)} \binom{n+l(\lambda)}{n} u_\lambda h_\lambda$$

where

$$(29) \quad u_\lambda = \frac{l(\lambda)}{\prod_{i \geq 1} m_i(\lambda)!}$$

Let (g_λ) be the adjoint basis of (h_λ^*) , that is

$$(30) \quad \langle g_\lambda, h_\mu^* \rangle = \delta_{\lambda\mu}$$

One has

$$(31) \quad g_n = -m_n = -p_n$$

The algebra morphism ψ^T (adjoint of ψ) is also an involution, and it maps g_λ to m_λ . The matrix Ψ of ψ is *strictly upper triangular* in the basis (h_λ) and one has

$$(32) \quad \Psi_{\lambda\lambda} = (-1)^{l(\lambda)}$$

Theorem 2.1 (Macdonald [10]). *The linear map*

$$(33) \quad \Phi : \begin{cases} R & \rightarrow \Lambda \\ c_\lambda & \mapsto g_\lambda \end{cases}$$

is an isomorphism of algebras.

2.4. Top connection coefficients. Now, let us recall some properties of the top connection coefficients $a_{\lambda\mu}^\nu$, which are the structure constants of the Farahat-Higman algebra R . By Theorem 2.1,

$$(34) \quad g_\lambda g_\mu = \sum_{|\nu|=|\lambda|+|\mu|} a_{\lambda\mu}^\nu g_\nu$$

When μ and ν are partitions of length 1 ($\mu = (r)$ and $\nu = (m)$), Macdonald gives in [10, ex. 24 p 131] an explicit formula for $a_{\lambda(r)}^{(m)}$, $m = |\lambda| + r$, that is

$$(35) \quad a_{\lambda(r)}^{(m)} = \begin{cases} \frac{(m+1)r!}{(r+1-l(\lambda)) \prod_{i>0} m_i(\lambda)!} & \text{if } l(\lambda) \leq r+1 \\ 0 & \text{otherwise} \end{cases}$$

Moreover, Macdonald gives a recurrence formula, for any partition ν with $|\nu| = |\lambda| + r$

$$(36) \quad a_{\lambda(r)}^\nu = \sum_{(i,\mu)/\mu \cup \nu = \lambda \cup (\nu_i)} a_{\mu(r)}^{(\nu_i)}$$

One can deduce from these formulas that the $a_{\lambda(r)}^\nu$ are zero except if $\nu \geq \lambda \cup (r)$, and that $a_{\lambda(r)}^{\lambda \cup (r)} > 0$. The multiplicative structure of the Farahat-Higman algebra is uniquely determined by (35) and (36).

2.5. Jucys-Murphy elements. Let n and i be two integers, $i \leq n$. Define

$$(37) \quad \xi_i = \sum_{j < i} (j, i) \in \mathbb{C}[\mathfrak{S}_n],$$

called the *ith Jucys-Murphy element*. It is the sum of all transpositions (i, j) with $j < i$. The Jucys-Murphy elements do not belong to the center Z_n of $\mathbb{C}[\mathfrak{S}_n]$, but they generate a maximal commutative subalgebra of $\mathbb{C}[\mathfrak{S}_n]$. Let

$$(38) \quad \Xi_n = \{\xi_1, \xi_2, \dots, \xi_n\}$$

be the alphabet of Jucys-Murphy elements. It is known that the algebra of symmetric functions in Ξ_n is exactly Z_n [8]. Hence, since $(C_\mu)_{\mu \vdash n}$ is a basis of Z_n , one can rewrite $f(\Xi_n)$ as

$$(39) \quad f(\Xi_n) = \sum_{\mu \vdash n} k_{f,\mu}(n) C_\mu$$

for any homogeneous symmetric function f , and this decomposition is unique. We can rewrite this formula in terms of the reduced cycle types

$$(40) \quad f(\Xi_n) = \sum c_{f,\mu}(n) c_\mu(n).$$

Let r be the degree of f . When $|\mu| = r$, $k_{f,\mu}(n)$ does not depend on n . Let us rewrite it as $k_{f,\mu}$ and define a new element $f(\Xi)$ of the Farahat-Higman algebra R , by considering only maximal terms in the expansion of $f(\Xi_n)$ for n large enough.

$$(41) \quad f(\Xi) = \sum_{\mu \vdash r} k_{f,\mu} c_\mu$$

Murray [12] has shown that for any f , the isomorphism Φ between R and Λ maps $f(\Xi)$ to $f(-X) = S_0(f)$.

$$(42) \quad \Phi(f(\Xi)) = f(-X)$$

Since $\Phi(c_\mu) = g_\mu$, using the duality between the bases (h_λ^\star) and (g_λ) , one has

$$(43) \quad k_{f,\mu} = \langle f(-X), h_\mu^\star \rangle$$

3. MORE ABOUT MURRAY'S RESULT

3.1. A new derivation. Let us give a new proof of (42). Lascoux and Thibon [9] give the formula

$$(44) \quad p_m(\Xi_n) = \sum_{k=1}^{m+1} \sum_{\kappa \vdash k, l(\kappa) \leq m-k+2} \phi_{\kappa,m} a_{\kappa,n}$$

where $a_{\kappa,n}$ is defined by

$$(45) \quad a_{\kappa,n} = \frac{1}{(n-k)!} z_{\kappa \cup 1^{n-k}} C_{\kappa \cup 1^{n-k}}$$

The C_λ are the classical conjugacy classes, and the $\phi_{\kappa,m}$ are defined from

$$(46) \quad \phi_\kappa(t) = \frac{(1-q^{-1})^{k-1} p_\kappa(q-1)}{k! z_k} \bigg|_{q=e^t}$$

by

$$(47) \quad \phi_\kappa(t) = \sum_{m \geq |\kappa| + l(\kappa) - 2} \phi_{\kappa,m} \frac{t^m}{m!}.$$

From (44), $C_{(m+1) \cup 1^{n-k}}$ is the only class of modified weight m which can give a contribution to $p_m(\Xi_n)$. Indeed, for $|\kappa| - l(\kappa) = m$, if we had $l(\kappa) > 1$ we would have $|\kappa| > m+1$, but the κ satisfying this inequality do not occur in the sum (44), so that the formula

$$(48) \quad p_m(\Xi) = \sum_{\mu \vdash m} k_\mu c_\mu$$

becomes

$$(49) \quad p_m(\Xi) = k_m c_m$$

Now we only have to determine the coefficient k_m . In order to do that, suppose that $n = m+1$. In this case we have

$$(50) \quad p_m(\Xi_{m+1}) = \xi_1^m + \xi_2^m + \dots + \xi_{m+1}^m,$$

that is,

$$(51) \quad p_m(\Xi_{m+1}) = 0^m + (1, 2)^m + ((1, 3) + (2, 3))^m + \dots \\ + ((1, m+1) + (2, m+1) + \dots + (m, m+1))^m$$

Only the last term can give a contribution to C_{m+1} . This term can be rewritten

$$(52) \quad \sum (i_1, m+1)(i_2, m+1) \dots (i_m, m+1)$$

where the sum is over the (i_1, i_2, \dots, i_m) where the i_k are integers such that $1 \leq i_k \leq m$. In this new sum, only terms with the i_k all distinct can give a contribution to C_{m+1} . The number of these terms is $m!$, that is, the cardinality of C_{m+1} , so that the coefficient of C_{m+1} in $p_m(\Xi_{m+1})$ is 1. This coefficient does not depend on n since $(m+1)$ has a maximal modified weight, so that the coefficient of c_m in $p_m(\Xi)$, that is k_m , is again 1, so that

$$(53) \quad p_m(\Xi) = c_m.$$

Hence, $\Phi(p_m(\Xi)) = \Phi(c_m) = g_m = -p_m$, and

$$(54) \quad \Phi(p_\lambda(\Xi)) = \Phi\left(\prod_i p_{\lambda_i}(\Xi)\right) = \prod_i \Phi(p_{\lambda_i}(\Xi)).$$

From that we have $\Phi(p_\lambda(\Xi)) = \prod_i (-p_{\lambda_i}) = (-1)^{l(\lambda)} p_\lambda = p_\lambda(-X)$, and since Λ is spanned by the p_μ , this implies (42).

3.2. Examples. Using his result and simple calculations in Λ , Murray [12] computes coefficients in the expansion of certain symmetric functions of the Jucys-Murphy elements over the c_λ . For example, he shows the following formulas

$$(55) \quad \langle e_k(-X), h_\lambda^* \rangle = 1$$

$$(56) \quad \langle h_k(-X), h_\lambda^* \rangle = \prod_i \text{Cat}_{\lambda_i-1}$$

(where Cat_i is the i th Catalan number) giving the top coefficients in the expansion of $e_k(\Xi)$ and $h_k(\Xi)$. In the same vein, let us give a new proof of a result of Matsumoto and Novak for the monomial functions. Let k be an integer, $\lambda \vdash k$ and $\mu \vdash k$. We denote by L_μ^λ the coefficient defined by

$$(57) \quad m_\lambda(\Xi) = \sum_{|\mu|=|\lambda|} L_\mu^\lambda c_\mu$$

Matsumoto and Novak [11] show that

$$(58) \quad L_\mu^\lambda = \sum_{(\lambda^{(1)}, \lambda^{(2)}, \dots) \in \mathfrak{R}(\lambda, \mu)} RC(\lambda^{(1)}) RC(\lambda^{(2)}) \dots$$

where

$$(59) \quad \mathfrak{R}(\lambda, \mu) = \{(\lambda^{(1)}, \lambda^{(2)}, \dots) / \forall i, \lambda^{(i)} \vdash \mu_i \text{ and } \lambda = \lambda^{(1)} \cup \lambda^{(2)} \cup \dots\}$$

and for any partition λ of N ,

$$(60) \quad RC(\lambda) = \frac{1}{N+1} m_\lambda(N+1) = \frac{|\lambda|!}{(|\lambda| - l(\lambda) + 1)! \prod_{i \geq 1} m_i(\lambda)!}$$

Matsumoto and Novak also give a combinatorial interpretation for $RC(\lambda)$. Let us show how to derive (58) from (43). We have

$$(61) \quad \begin{aligned} L_\mu^\lambda &= \langle m_\lambda(-X), h_\mu^* \rangle \\ &= \langle m_\lambda, h_\mu^*(-X) \rangle \\ &= \left\langle m_\lambda, \frac{h_{\mu_1}((\mu_1+1)X) h_{\mu_2}((\mu_2+1)X) \dots}{(\mu_1+1)(\mu_2+1) \dots} \right\rangle \\ &= \frac{1}{\prod_i (\mu_i+1)} \left\langle m_\lambda, \prod_i \sum_{\lambda^{(i)} \vdash \mu_i} m_{\lambda^{(i)}}(\mu_i+1) h_{\lambda^{(i)}} \right\rangle \end{aligned}$$

Expanding the right factor of the scalar product, we obtain nonzero terms only if $h_{\lambda^{(1)}} h_{\lambda^{(2)}} \dots = h_\lambda$, since otherwise one has $\langle m_\lambda, h_{\lambda^{(1)}} h_{\lambda^{(2)}} \dots \rangle = 0$. Hence,

$$(62) \quad L_\mu^\lambda = \frac{1}{\prod_i (\mu_i+1)} \sum (m_{\lambda^{(1)}}(\mu_1+1)) (m_{\lambda^{(2)}}(\mu_2+1)) \dots \langle m_\lambda, h_\lambda \rangle$$

The sum is over the $(\lambda^{(1)}, \lambda^{(2)}, \dots)$ such that $\lambda^{(i)} \vdash \mu_i$ for all i and $\bigcup_i \lambda^{(i)} = \lambda$, that is, over the set $\mathfrak{R}(\lambda, \mu)$. Moreover, one has $\langle m_\lambda, h_\lambda \rangle = 1$, so that

$$(63) \quad L_\mu^\lambda = \sum_{(\lambda^{(1)}, \lambda^{(2)}, \dots) \in \mathfrak{R}(\lambda, \mu)} \frac{m_{\lambda^{(1)}}(\mu_1+1)}{\mu_1+1} \frac{m_{\lambda^{(2)}}(\mu_2+1)}{\mu_2+1} \dots$$

This is the result of Matsumoto and Novak.

3.3. Coefficient of the cycle with maximal length in $p_\lambda(\Xi_n)$. We call *modified cycle type* of a product of cycles the sequence $(l_1 - 1, l_2 - 1, \dots)$, where the l_i are the lengths of the factors. For example, the product (13)(234) has modified cycle type $(2 - 1, 3 - 1) = (1, 2)$. Biane [1] obtains an explicit formula for the number α_λ of factorisations with modified cycle type λ for a cycle of length $n + 1$, where $n = |\lambda| = \sum_i \lambda_i$, that is

$$(64) \quad \alpha_\lambda = (n+1)^{l(\lambda)-1}$$

Let us prove this with symmetric functions. We consider in the Farahat-Higman algebra R the product

$$(65) \quad c^\lambda = c_{\lambda_1} c_{\lambda_2} \dots$$

Since R is isomorphic to Λ , it is commutative, so that the order of the elements of λ has no importance. Hence, we can assume that λ is a partition. One has

$$(66) \quad \begin{aligned} \Phi(c^\lambda) &= g_{\lambda_1} g_{\lambda_2} \dots = (-p_{\lambda_1})(-p_{\lambda_2}) \dots \\ &= (-1)^{l(\lambda)} p_\lambda = p_\lambda(-X) \end{aligned}$$

Hence, from (42),

$$(67) \quad c^\lambda = p_\lambda(\Xi)$$

Moreover, c^λ is the sum of all cycle products of modified type λ , and c_n is the sum of all the cycles of length $n + 1$. Hence, the total number of cycle products of modified type λ is $k_\lambda^n \text{card}(c_n)$, where k_λ^n is the coefficient of c_n in c^λ . The number α_λ of these factorisations does not depend on the choice of the cycle, so that $\alpha_\lambda = k_n^\lambda \frac{\text{card}(c_n)}{\text{card}(c_n)} = k_\lambda^n$, which is also from (67) the coefficient of c_n in $p_\lambda(\Xi)$. Hence, one has

$$(68) \quad \alpha_\lambda = (-1)^{l(\lambda)} \langle p_\lambda, h_n^* \rangle,$$

so that

$$(69) \quad \alpha_\lambda = \frac{1}{n+1} (n+1)^{l(\lambda)} \langle p_\lambda, h_n \rangle = (n+1)^{l(\lambda)-1} \langle p_\lambda, h_n \rangle$$

(since the bases (m_λ) and (h_λ) are dual to each other, $\langle p_\lambda, h_n \rangle$ is the coefficient of m_n in p_λ , that is, 1).

3.4. Generalization. Here, we give a combinatorial interpretation for the coefficient of c_μ in $p_\lambda(\Xi)$. Now we have

$$(70) \quad \langle p_\lambda, h_n^* \rangle = (-1)^{l(\lambda)} (n+1)^{l(\lambda)-1}$$

On another hand, since $p_\lambda(\Xi) = c_{\lambda_1} c_{\lambda_2} \dots$ in R ,

$$(71) \quad c_{\lambda_1} c_{\lambda_2} \dots = \sum_{|\lambda|=|\mu|} \langle p_\lambda(-X), h_\mu^* \rangle c_\mu,$$

so that $\langle p_\lambda(-X), h_\mu^* \rangle$ corresponds to the total number of decompositions in an arbitrary product of modified type λ of a product of disjoint cycles with modified type μ . In such a decomposition, each c_{μ_i} comes from certain $\prod_i c_{\lambda^{(i)}}$, where $\lambda^{(i)} \vdash \mu_i$ is a subpartition of λ , with $\lambda^{(1)} \cup \lambda^{(2)} \cup \dots = \lambda$, that is, $(\lambda^{(1)}, \lambda^{(2)}, \dots) \in \mathfrak{R}(\lambda, \mu)$. Moreover, each $(\lambda^{(1)}, \lambda^{(2)}, \dots)$ must be counted $m(\lambda^{(1)}, \lambda^{(2)}, \dots)$ times, where

$$(72) \quad m(\lambda^{(1)}, \lambda^{(2)}, \dots) = \prod_{j \geq 1} \frac{m_j(\lambda)!}{m_j(\lambda^{(1)})! m_j(\lambda^{(2)})! \dots}$$

For a given c_{μ_i} , the number of decompositions of c_{μ_i} of a certain type $\prod_i c_{\lambda^{(i)}}$ corresponds to the coefficient of c_{μ_i} in $p_{\lambda^{(i)}}(\Xi)$, that is $\langle p_{\lambda^{(i)}}(-X), h_{\mu_i}^* \rangle$, so that we have from (70)

$$(73) \quad \langle p_\lambda(-X), h_\mu^* \rangle = \sum_{(\lambda^{(1)}, \lambda^{(2)}, \dots) \in \mathfrak{R}(\lambda, \mu)} m(\lambda^{(1)}, \lambda^{(2)}, \dots) \prod_i (\mu_i + 1)^{l(\lambda^{(i)})-1}$$

Note that it is nonzero only if λ is a refinement of μ . Finally, one has

$$(74) \quad p_\lambda(\Xi) = \sum_{|\lambda|=|\mu|} \left(\sum_{(\lambda^{(1)}, \lambda^{(2)}, \dots) \in \mathfrak{R}(\lambda, \mu)} \left(\prod_j \frac{m_j(\lambda)!}{m_j(\lambda^{(1)})! m_j(\lambda^{(2)})! \dots} \right) \left(\prod_i (\mu_i + 1)^{l(\lambda^{(i)})-1} \right) \right) c_\mu$$

Note that from (70) we have

$$(75) \quad h_n^* = \sum_{\mu \vdash n} (-1)^{l(\mu)} (n+1)^{l(\mu)-1} \frac{p_\mu}{z_\mu}$$

Expanding h_μ^* by means of this expression, since we have for $\bigcup_i \lambda^{(i)} = \lambda$

$$(76) \quad \prod_{j \geq 1} \frac{m_j(\lambda)!}{m_j(\lambda^{(1)})! m_j(\lambda^{(2)})! \dots} = \frac{z_\lambda}{z_{\lambda^{(1)}} z_{\lambda^{(2)}} \dots}$$

we see that (73) can also be obtained by simple calculations in Λ .

4. THE FAÀ DI BRUNO ALGEBRA AND ITS DEFORMATION

4.1. The Faà di Bruno algebra. There is another coproduct on Λ , denoted here by Δ_1 and defined by

$$(77) \quad \Delta_1 h_n = \sum_{k=0}^n h_k(X) \otimes h_{n-k}((k+1)X)$$

or equivalently

$$(78) \quad \Delta_1 h_n = \sum_{k=0}^n \sum_{\mu \vdash n-k} m_\mu(k+1) h_k \otimes h_\mu$$

This coproduct defines a Hopf algebra with the counit ϵ as in \mathcal{H}_0 , and the antipode $S_1 = \psi$, that is, the involution mapping h_λ to h_λ^* . The Hopf algebra

$$(79) \quad \mathcal{H}_1 = (\Lambda, \cdot, 1, \Delta_1, \epsilon, S_1)$$

is called the *Faà di Bruno algebra*. It has the following interpretation. Let

$$(80) \quad G_1 = \{\alpha \mid \exists a \in G_0, \forall t, \alpha(t) = ta(t)\} = tG_0 = t + t^2\mathbb{C}[[t]]$$

be the group of formal diffeomorphisms of the line tangent to the identity, and let again k_n be the linear form

$$(81) \quad k_n : \alpha(t) = t + a_1 t^2 + a_2 t^3 + \dots \mapsto a_n$$

Let Δ be the coproduct such that $\Delta k_n(\alpha, \beta)$ is the coefficient of t^{n+1} in

$$(82) \quad (\alpha \circ \beta)(t) = \alpha(\beta(t))$$

The bialgebra \mathcal{F} defined by this coproduct is isomorphic to \mathcal{H}_1 under the correspondence $k_n \mapsto h_n$.

4.2. A deformation of the Faà di Bruno algebra. Let γ be a real parameter in $[0, 1]$, and Δ_γ be the coproduct on Λ defined by

$$(83) \quad \Delta_\gamma(h_n) = \sum_{k=0}^n h_k \otimes h_{n-k}((k\gamma + 1)X)$$

or equivalently

$$(84) \quad \Delta_\gamma(h_n) = \sum_{k=0}^n \sum_{\mu \vdash n-k} m_\mu(k\gamma + 1) h_k \otimes h_\mu$$

Foissy obtains this coproduct in [5] in his investigation of formal Dyson-Schwinger equations in the Connes-Kreimer Hopf algebra, and shows that for $\gamma \in]0, 1]$, the

resulting bialgebras \mathcal{H}_γ are Hopf algebras, isomorphic to the Faà di Bruno algebra. Obviously, Δ_0 corresponds to \mathcal{H}_0 .

4.3. Graded dual of the deformed Faà di Bruno algebra. Consider now the Hopf algebra \mathcal{H}'_γ , the graded dual of the deformation \mathcal{H}_γ considered in section 3. Denote by \star_γ the product on \mathcal{H}'_γ . For any $f \in \mathcal{H}'_\gamma$ and any symmetric function g in \mathcal{H}_γ , denote by $\langle f, g \rangle$ the action of f on g . One has for any $f \in \mathcal{H}'_\gamma$, $g \in \mathcal{H}'_\gamma$ and $h \in \Lambda$, by definition of a dual Hopf algebra,

$$(85) \quad \langle f \star_\gamma g, h \rangle = \langle f \otimes g, \Delta_\gamma(h) \rangle$$

where

$$(86) \quad \langle a \otimes b, c \otimes d \rangle = \langle a, c \rangle \langle b, d \rangle$$

Denote by (d_λ) , (q_λ) and (b_λ) the bases respectively dual to (h_λ) , $\left(\frac{p_\lambda}{z_\lambda}\right)$ and h_λ^* in the sense of \mathcal{H}'_γ , that is, for example

$$(87) \quad \langle d_\lambda, h_\mu \rangle = \delta_{\lambda\mu}$$

For any γ in $[0, 1]$,

$$(88) \quad q_n = d_n = -b_n$$

Note that these b_n generate \mathcal{H}'_γ . \mathcal{H}_0 is *self-dual*, so that in the case $\gamma = 0$,

$$(89) \quad d_\lambda = m_\lambda, \quad q_\lambda = p_\lambda, \quad b_\lambda = g_\lambda$$

When $\gamma \neq 0$, \mathcal{H}_γ is not self-dual and not commutative, since the coproduct Δ_γ on Λ is not cocommutative.

4.4. Multiplicative structure of \mathcal{H}'_γ . Now, let f and g be two elements of \mathcal{H}'_γ . One has for any partition μ ,

$$\begin{aligned} \langle f \star_\gamma g, h_\mu \rangle &= \langle f \otimes g, \Delta_\gamma(h_\mu) \rangle \\ &= \langle f \otimes g, \Delta_\gamma(h_{\mu_1}) \Delta_\gamma(h_{\mu_2}) \dots \rangle \\ &= \left\langle f \otimes g, \prod_i \sum_{k_i + l_i = \mu_i} h_{k_i} \otimes h_{l_i} ((\gamma k_i + 1)X) \right\rangle \\ &= \left\langle f \otimes g, \sum_{k_i + l_i = \mu_i} \prod_i h_{k_i} \otimes h_{l_i} ((\gamma k_i + 1)X) \right\rangle \\ (90) \quad &= \sum \langle f \otimes g, \prod_i h_{k_i} \otimes h_{l_i} ((\gamma k_i + 1)X) \rangle \end{aligned}$$

where the parameters of the sum are the same as above. Hence,

$$\begin{aligned} (91) \quad \langle f \star_\gamma g, h_\mu \rangle &= \sum \langle f, h_{(k_1, k_2, \dots)} \rangle \langle g, \prod_i h_{l_i} (\gamma k_i + 1)X \rangle \\ &= \sum \langle f, h_{(k_1, k_2, \dots)} \rangle \langle g, \prod_i \sum_{\rho \vdash l_i} m_\rho (\gamma k_i + 1) h_\rho \rangle \end{aligned}$$

Now let us change the parameters of the sum, considering the partitions ρ_i , with $|\rho_i| = l_i$ and $k_i = \mu_i - |\rho_i|$. The sum is now over the $(\rho_1, \rho_2, \dots, \rho)$, such that for all i , $|\rho_i| \leq \mu_i$, and ρ is the union of the ρ_i . We obtain

$$(92) \quad \langle f \star_\gamma g, h_\mu \rangle = \sum \langle f, h_{(\mu_1-|\rho_1|, \mu_2-|\rho_2|, \dots)} \rangle \langle g, \prod_i m_{\rho_i}(\gamma\mu_i - \gamma|\rho_i| + 1) h_{\rho_i} \rangle,$$

and finally,

$$(93) \quad \langle f \star_\gamma g, h_\mu \rangle = \sum \left(\prod_i m_{\rho_i}(\gamma\mu_i - \gamma|\rho_i| + 1) \right) \langle f, h_{(\mu_1-|\rho_1|, \mu_2-|\rho_2|, \dots)} \rangle \langle g, h_\rho \rangle$$

One can use this formula to expand $f \star_\gamma g$ on the d_μ , where f and g are any two elements in \mathcal{H}'_γ

4.5. Action of q_n^\perp on the h_μ . We define an operator q_n^\perp on Λ as follows, for $f \in \mathcal{H}'_\gamma$ and $g \in \mathcal{H}_\gamma$.

$$(94) \quad \langle f \star_\gamma q_n, g \rangle = \langle f, q_n^\perp g \rangle$$

The operator q_n^\perp is a *derivation*, because it is the adjoint of the right multiplication by a *primitive element*:

$$(95) \quad q_n^\perp(fg) = f q_n^\perp(g) + q_n^\perp(f)g$$

We shall need its action on the h_μ . Let $n > 0$ and μ be a partition. We can write from (93) :

$$(96) \quad \langle f \star_\gamma q_n, h_\mu \rangle = \sum \left(\prod_i m_{\rho_i}(\gamma\mu_i - \gamma|\rho_i| + 1) \right) \langle f, h_{(\mu_1-|\rho_1|, \mu_2-|\rho_2|, \dots)} \rangle \langle q_n, h_\rho \rangle$$

Since $q_n = d_n$ for all n , one has $\langle q_n, h_\rho \rangle = \delta_{n\rho}$, so that a term in this sum gives a nonzero contribution only if

$$(97) \quad \rho = (n)$$

In this case we have also $\langle q_n, h_\rho \rangle = 1$, and

$$(98) \quad \langle f \star_\gamma q_n, h_\mu \rangle = \sum_{i/\mu_i \geq n} m_n(\gamma\mu_i - \gamma n + 1) \langle f, h_{\mu \setminus \mu_i \cup (\mu_i - n)} \rangle$$

Since $m_n(\gamma\mu_i - \gamma n + 1) = \gamma(\mu_i - n) + 1$, one can deduce

$$(99) \quad q_n^\perp h_\mu = \sum_{i/\mu_i \geq n} (\gamma\mu_i - \gamma n + 1) h_{\mu \setminus \mu_i \cup (\mu_i - n)}$$

When μ consists only in one part N , this formula can be rewritten

$$(100) \quad q_n^\perp h_N = (\gamma N - \gamma n + 1) h_{N-n}$$

Since q_n^\perp is a derivation, one has its action on the multiplicative basis (h_μ) , and one can use (100) to fully explicit it. We have $q_n^\perp = D_n + E_n$, where D_n and E_n are the derivations defined by

$$(101) \quad D_n h_n = h_{N-n}$$

and

$$(102) \quad E_n h_n = \gamma(N - n)h_{N-n},$$

that is,

$$(103) \quad D_n = \sum_{r \geq 0} h_r \frac{\partial}{\partial h_{n+r}}$$

and

$$(104) \quad E_n = \gamma \sum_{r \geq 0} r h_r \frac{\partial}{\partial h_{n+r}}$$

so that

$$(105) \quad q_n^\perp = \sum_{r \geq 0} (1 + \gamma r) h_r \frac{\partial}{\partial h_{n+r}}$$

We can see that this is valid also in the degenerate case $\gamma = 0$.

4.6. Action of q_n^\perp on the p_μ . From (100), we have

$$(106) \quad q_n^\perp h_N = p_n^\perp h_N + \gamma(N - n)h_{N-n}$$

On another hand,

$$(107) \quad (N - n)h_{N-n} = p_1 h_{N-n-1} + p_2 h_{N-n-2} + p_3 h_{N-n-3} + \dots$$

Then,

$$(108) \quad \begin{aligned} q_n^\perp h_N &= p_n^\perp h_N + \gamma p_1 h_{N-n-1} + \gamma p_2 h_{N-n-2} + \dots \\ &= p_n^\perp h_N + \gamma p_1 (p_{n+1}^\perp h_N) + \gamma p_2 (p_{n+2}^\perp h_N) + \dots \\ &= p_n^\perp h_N + \gamma \sum_{r > n} p_{r-n} p_r^\perp h_N \end{aligned}$$

so that

$$(109) \quad q_n^\perp = p_n^\perp + \gamma \sum_{r > n} p_{r-n} p_r^\perp$$

5. A DEFORMATION OF THE FARAHAT-HIGMAN ALGEBRA

5.1. Recurrences for the structure constants of \mathcal{H}'_γ . Denote by $a_{\lambda,\mu}^\nu(\gamma)$ the structure constants in the basis b_μ .

$$(110) \quad b_\lambda \star_\gamma b_\mu = \sum_\nu a_{\lambda,\mu}^\nu(\gamma) b_\nu$$

When $\gamma = 0$, the b_μ are identified with the g_μ , and then $a_{\lambda,\mu}^\nu(0)$ coincides with $a_{\lambda,\mu}^\nu$, the top connection coefficient.

Since $b_n = -q_n$, one has

$$(111) \quad a_{\lambda,(n)}^\nu(\gamma) = \langle b_\lambda \star_\gamma b_n, h_\nu^* \rangle = -\langle b_\lambda \star_\gamma q_n, h_\nu^* \rangle$$

Hence,

$$(112) \quad a_{\lambda,(n)}^\nu(\gamma) = -\langle b_\lambda, q_n^\perp h_\nu^* \rangle$$

Since q_n^\perp is a derivation, one can rewrite this as

$$(113) \quad a_{\lambda,(n)}^\nu(\gamma) = -\sum_i \langle b_\lambda, h_{\nu \setminus (\nu_i)}^* q_n^\perp(h_{\nu_i}^*) \rangle$$

The i th term in this sum corresponds to the coefficient of h_λ^* in

$$(114) \quad h_{\nu \setminus (\nu_i)}^* q_n^\perp(h_{\nu_i}^*)$$

Hence this term gives a nonnegative contribution only if there exists a partition μ such that

$$(115) \quad (\nu \setminus (\nu_i)) \cup \mu = \lambda,$$

that is,

$$(116) \quad \mu \cup \nu = \lambda \cup (\nu_i)$$

Hence, the i th term is also equal to the coefficient of h_μ^* in $q_n^\perp(h_{\nu_i}^*)$, that is, $a_{\mu(n)}^{\nu_i}(\gamma)$. Summarizing, we have proved:

Theorem 5.1. *The structure constants $a_{\lambda,(n)}^\nu(\gamma)$ satisfy the recursion*

$$(117) \quad a_{\lambda,(n)}^\nu(\gamma) = \sum a_{\mu(n)}^{\nu_i}(\gamma)$$

where the sum is over the (i, μ) such that

$$(118) \quad \mu \cup \nu = \lambda \cup (\nu_i)$$

This formula is a generalization of (36), which is recovered for $\gamma = 0$.

5.2. Multiplicative structure of the deformed Farahat-Higman algebra. In the case where ν has only one part, there is a closed formula for $a_{\lambda(r)}^\nu(\gamma)$.

Theorem 5.2. *For $r \in \mathbb{N}$, $N \in \mathbb{N}$ and μ a partition,*

$$(119) \quad a_{\lambda,(r)}^{(N)}(\gamma) = \left(1 - \frac{N-r}{N+1}\gamma\right) a_{\lambda,(r)}^{(N)}(0)$$

$a_{\lambda,(r)}^{(N)}(0)$ corresponds to $a_{\lambda,(r)}^{(N)}$ in formula (35), where m is replaced by N .

Together with the recurrence formula (117), this determines completely the multiplicative structure of \mathcal{H}'_γ , since it is generated by the b_n . In order to derive (119), we shall need the following lemmas.

Lemma 5.3. *Let r and n be two positive integers, with $r < n$. Then,*

$$(120) \quad p_r^\perp h_n^* = \sum_{\rho \vdash n-r} (-1)^{l(\rho)-1} (n+1)^{l(\rho)} \frac{p_\rho}{z_\rho}$$

Proof. From (75) we have

$$(121) \quad p_r^\perp h_n^\star = \sum_{\mu \vdash n} \frac{(-1)^{l(\mu)} (n+1)^{l(\mu)-1}}{z_\mu} p_r^\perp(p_\mu)$$

The action of p_r^\perp on p_μ is

$$(122) \quad p_r^\perp(p_\mu) = r \frac{\partial p_\mu}{\partial p_r} = r m_r(\mu) p_{\mu \setminus (r)}$$

On another hand,

$$(123) \quad z_{\mu \setminus (r)} = \frac{z_\mu m_r(\mu \setminus (r))!}{r m_r(\mu)!}$$

with $m_r(\mu \setminus (r)) = m_r(\mu) - 1$, hence $\frac{m_r(\mu)!}{m_r(\mu \setminus (r))!} = m_r(\mu)$, so that

$$(124) \quad \frac{z_\mu}{r m_r(\mu)} = z_{\mu \setminus (r)}$$

Hence,

$$(125) \quad \frac{p_r^\perp(p_\mu)}{z_\mu} = \frac{p_{\mu \setminus (r)}}{z_{\mu \setminus (r)}}$$

and

$$(126) \quad p_r^\perp h_n^\star = \sum_{\mu \vdash n} (-1)^{l(\mu)} (n+1)^{l(\mu)-1} \frac{p_{\mu \setminus (r)}}{z_{\mu \setminus (r)}}$$

or, equivalently,

$$(127) \quad p_r^\perp h_n^\star = \sum_{\rho \vdash n-r} (-1)^{l(\rho \cup (r))} (n+1)^{l(\rho \cup (r))-1} \frac{p_\rho}{z_\rho}$$

Since $l(\rho \cup (r)) = l(\rho) + 1$, we deduce the required formula . ■

Lemma 5.4. *Let r and N be two positive integers, with $N > r$. Then,*

$$(128) \quad \sum_{n > r} p_{n-r} (p_n^\perp h_N^\star) = -\frac{N-r}{N+1} p_r^\perp h_N^\star$$

Proof. From (120), one has

$$(129) \quad \sum_{n > r} p_{n-r} (p_n^\perp h_N^\star) = \sum_{n > r} \sum_{\rho \vdash n-r} (-1)^{l(\rho)+1} (n+1)^{l(\rho)} \frac{p_{\rho \cup (n-r)}}{z_\rho}$$

for $\mu = \rho \cup (n-r)$ and $k = n-r$, we obtain :

$$(130) \quad \sum_{n > r} p_{n-r} (p_n^\perp h_N^\star) = \sum_{\mu \vdash N-r} \sum_{k \in \mu} (-1)^{l(\mu)} (N+1)^{l(\mu)-1} \frac{p_\mu}{z_{\mu \setminus (k)}}$$

$$(131) \quad \sum_{n > r} p_{n-r} (p_n^\perp h_N^\star) = \sum_{\mu \vdash N-r} (-1)^{l(\mu)} (N+1)^{l(\mu)-1} \left(\sum_{k \in \mu} \frac{1}{z_{\mu \setminus (k)}} \right) p_\mu$$

From (124), one has

$$(132) \quad \sum_{k \in \mu} \frac{1}{z_{\mu \setminus (k)}} = \sum_{k \in \mu} \frac{km_{\mu}(k)}{z_{\mu}}$$

Hence,

$$(133) \quad \sum_{k \in \mu} \frac{1}{z_{\mu \setminus (k)}} = \frac{1}{z_{\mu}} \sum_{k \in \mu} km_{\mu}(k) = \frac{|\mu|}{z_{\mu}}$$

So we can rewrite (131) as

$$(134) \quad \sum_{n > r} p_{n-r}(p_n^{\perp} h_N^{\star}) = \sum_{\mu \vdash N-r} (-1)^{l(\mu)} (N+1)^{l(\mu)-1} \frac{N-r}{z_{\mu}} p_{\mu}$$

$$(135) \quad \sum_{n > r} p_{n-r}(p_n^{\perp} h_N^{\star}) = \frac{N-r}{N+1} \sum_{\mu \vdash N-r} (-1)^{l(\mu)} (N+1)^{l(\mu)} \frac{p_{\mu}}{z_{\mu}}$$

From (120), we get the required formula. ■

Proof – (of Theorem 5.2) Since $b_r = -q_r$ and $g_r = -p_r$, one has

$$(136) \quad \langle b_{\lambda} \star_{\gamma} b_r, h_N^{\star} \rangle = -\langle b_{\lambda}, q_r^{\perp} h_N^{\star} \rangle$$

Then, from (109) and lemma 5.3,

$$(137) \quad \begin{aligned} \langle b_{\lambda} \star_{\gamma} b_r, h_N^{\star} \rangle &= -\langle b_{\lambda}, p_r^{\perp} h_N^{\star} \rangle - \gamma \langle b_{\lambda}, \sum_{n > r} p_{n-r}(p_n^{\perp} h_N^{\star}) \rangle \\ &= -\langle g_{\lambda}, p_r^{\perp} h_N^{\star} \rangle + \gamma \frac{N-r}{N+1} \langle b_{\lambda}, p_r^{\perp} h_N^{\star} \rangle \\ &= -\langle g_{\lambda} p_r, h_N^{\star} \rangle + \gamma \frac{N-r}{N+1} \langle g_{\lambda}, p_r^{\perp} h_N^{\star} \rangle \\ &= \langle g_{\lambda} g_r, h_N^{\star} \rangle - \gamma \frac{N-r}{N+1} \langle g_{\lambda} g_r, h_N^{\star} \rangle \\ &= \left(1 - \gamma \frac{N-r}{N+1} \right) \langle g_{\lambda} g_r, h_N^{\star} \rangle \end{aligned}$$

From that, we deduce (119).

6. A DEFORMATION OF THE WITT ALGEBRA

6.1. A simpler multiplicative formula in \mathcal{H}'_{γ} . Now let us expand $q_k \star_{\gamma} q_n$ on the q_{μ} . One has

$$(138) \quad q_k \star_{\gamma} q_n = \sum_{\mu} \frac{1}{z_{\mu}} \langle q_k \star_{\gamma} q_n, p_{\mu} \rangle q_{\mu},$$

so that

$$(139) \quad q_k \star_{\gamma} q_n = \sum_{\mu} \frac{1}{z_{\mu}} \langle q_k, q_n^{\perp}(p_{\mu}) \rangle q_{\mu}$$

From (109), deduce that

$$(140) \quad q_n^\perp p_\mu = p_n^\perp p_\mu + \gamma \sum_{r>n} p_{r-n} p_r^\perp p_\mu$$

so that

$$(141) \quad \begin{aligned} q_n^\perp p_\mu &= p_n^\perp p_\mu + \gamma \sum_{r>n} r p_{r-n} \frac{\partial p_\mu}{\partial p_r} \\ &= p_n^\perp p_\mu + \gamma \sum_{r>n} r m_r(\mu) p_{\mu \setminus (r) \cup (r-n)} \end{aligned}$$

Hence, q_μ gives to $q_k \star_\gamma q_n$ a contribution to the factor of γ only if $\mu = (k+n)$. In this case we have

$$(142) \quad \langle q_k, q_n^\perp p_{k+n} \rangle = \gamma(k+n)k$$

Hence,

$$(143) \quad q_k \star_\gamma q_n = q_{(k,n)} + k\gamma q_{k+n}$$

this formula also completely determines the multiplicative structure of \mathcal{H}'_γ . In the case $\gamma = 1$, we can see that it is coherent with the result of [3].

6.2. Lie algebra structure corresponding to \mathcal{H}'_γ . Now, suppose that $\gamma \neq 0$. Since \mathcal{H}'_γ is a connected cocommutative Hopf algebra, it is the universal enveloping algebra of a Lie algebra \mathcal{L}_γ . From (143), its bracket is determined by

$$(144) \quad [q_k, q_n]_\gamma = \gamma(k-n)q_{k+n}.$$

Denote by $d_{n,\gamma}$ the differential operator

$$(145) \quad d_{n,\gamma} = t^{1-n\gamma} \frac{d}{dt}$$

The $d_{n,\gamma}$ satisfy the same relation (144) as the q_n of \mathcal{L}_γ , so that \mathcal{H}'_γ can be interpreted as a Lie algebra of differential operators: it is the Lie algebra generated by the $d_{n,\gamma}$. In the case $\gamma = 1$ one has $d_{n,1} = t^{1-n} \frac{d}{dt}$. These operators generate \mathcal{L}_1 , that is called the *Witt algebra*. It is known that the universal enveloping algebra of the Witt algebra is the dual of the Fàa di Bruno algebra. Note also that from the commutativity of \mathcal{H}'_0 , (144) is also valid in the degenerate case $\gamma = 0$.

REFERENCES

- [1] P. BIANE, *Minimal factorizations of a Cycle and Central Multiplicative Functions on the Infinite Symmetric Group*, J. Combin. Theor. A **76** (1996), 97–212.
- [2] S. CORTEEL, A. GOUPIL and G. SCHAEFFER, *Content evaluation and class symmetric functions*, Adv. Math. **188** (2004), no. 2, 315–336.
- [3] H. FIGUEROA and J. M. GRACIA-BONDIA, *Combinatorial Hopf algebras in quantum field theory I*, Rev.Math.Phys. **17** (2005), 881–976.
- [4] H. K. FARAHAT and G. HIGMAN, *The centres of symmetric group rings*, Proc. Roy. Soc. London Ser. A **250** (1959), 212–221.
- [5] L. FOISSY, *Faà di Bruno subalgebras of the Hopf algebra of planar trees from combinatorial Dyson-Schwinger equations*, Adv. Math. **218** (2008), no. 1, 136–162.

- [6] I. P. GOULDEN and D. M. JACKSON, *Symmetric functions and Macdonald's result for top connexion coefficients in the symmetric group*, J. Algebra **166** (1994), no. 2, 364–378.
- [7] I. P. GOULDEN and D. M. JACKSON, *Connexion coefficients for the symmetric group, free products in operator algebras and random matrices*, in Fields Institute Communications: Free Probability Theory, D.-V. Voiculescu, Ed. Providence, RI: Amer. Math. Soc., 1997, vol. 12, pp. 105–125.
- [8] A.-A.A. JUCYS, *Symmetric polynomials and the center of the symmetric group ring*, Rep. Mathematical Phys. **5** (1974), no. 1, 107–112.
- [9] A. LASCOUX and J.-Y. THIBON, *Vertex operators and the class algebras of symmetric groups*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **283** (2001), Teor. Predst. Din. Sist. Komb. i Algoritm. Metody. 6, 156–177, 261; also in J. Math. Sci. (N. Y.) **121** (2004), no. 3, 2380–2392
- [10] I.G. MACDONALD, *Symmetric functions and Hall polynomials*, 2nd ed., Oxford University Press, 1995.
- [11] S. MATSUMOTO and J. NOVAK, *Jucys-Murphy Elements and Unitary Matrix Integrals*, arXiv:0905.1992v2.
- [12] J. MURRAY, *Generators for the centre of the group algebra of a symmetric group*, J. Algebra **271** (2004), 725–748.

INSTITUT GASPARD MONGE, UNIVERSITÉ PARIS-EST MARNE-LA-VALLÉE, 5 BOULEVARD DESCARTES,
CHAMPS-SUR-MARNE, 77454 MARNE-LA-VALLÉE CEDEX 2, FRANCE
E-mail address: bultel@univ-mlv.fr